

INFECTIONS

4. ANTIFUNGALS

GYNO-MIKOZAL

Presentation

Cream 2%: Collapsible tube of 78g of the cream with an applicator.

Vaginal ovules 200mg: Pack of 7 ovules.

Vaginal ovules 400mg: Pack of 3 ovules.

Composition

Each 1 gram of the cream contains: Miconazole nitrate 20mg

Each vaginal ovule contains: Miconazole nitrate 200mg or 400mg.

Indications

Indicated for the local treatment of vulvovaginal candidiasis caused by *Candida albicans* and other species of *Candida* in pregnant (second and third trimesters only) and nonpregnant women. It may also be used for the treatment of mycotic balanitis.

Since miconazole nitrate is effective against certain strains of Gram-positive bacteria, **Gyno-Mikozal** may be used in cases of mycoses secondarily infected with such bacteria.

Dosage and Administration

Cream 2%: The content of one applicator (5g) of cream should be emptied deeply into the vagina once daily at bedtime for 10 - 14 days or twice daily for 7 days.

Vaginal ovules: A single daily dose of 200mg vaginal ovule for one week or 400mg vaginal ovule for 3 days to be inserted deeply into the vagina at bedtime.

The entire course of treatment should be completed even when the symptoms disappear rapidly. It should be

continued during menstruation. If needed, a further course of treatment can be given.

For the treatment of vulvitis and other superficial sites of infection that are almost invariably associated with vaginal candidiasis, vaginal application of **Gyno-Mikozal** cream or ovules can be supplemented with topical application of the cream to anogenital area twice daily.

